

Vitamin C
orthomed.com

For MD's only

Preparation of Sodium Ascorbate for IV and IM Use

ROBERT F. CATHCART III, M.D.
ALLERGY, ENVIRONMENTAL & ORTHOMOLECULAR MEDICINE
127 SECOND STREET, SUITE 4
LOS ALTOS, CALIFORNIA 94022
(650) 949-2822
FAX (650) 949-5083

Note: some of these solutions are labeled ascorbic acid which is confusing. They are all buffered so as to really make them mostly sodium ascorbate.

Note: the following are excerpts from letters sent to physicians on the subject of IVC.

If one does not want to make their sodium ascorbate stock solutions from scratch like I recommend (and I can well understand why you might not): You can order from

Merit Pharmaceuticals, 2611 San
Fernando, Los Angeles, CA 90065,
For CA 800-696-3748
Out-of State 800-421-9657

[To order the Sodium Ascorbate Fine Crystals](#)

Wholesale Nutrition
915 S. San Tomas Equino Road
Campbell, CA 95008
order 800-325-2664 or FAX 408-867-6236

The Stock Bottle of Sodium Ascorbate

Sterilize a 500 cc IV bottle along with a funnel, the rubber stopper, and a spoon. Then fill the bottle to the 300 cc line with [sodium ascorbate fine crystals](#). (I weighed the sodium ascorbate out one time and 250 gm came up to the 300 cc line.) Then add 1/3 of the 20 ml bottle (6.6 cc) of edetate disodium injection, USP 150 mg/ml. Then add water for injection q.s. 500 cc. Shake up the bottle and if there is 1 mm of crystals left on the bottom, add 1 mm of water to the top. It turns out that sodium ascorbate is soluble to almost exactly a 50% concentration at room temperature. I do not worry about the sterility of this because this is very bacteriocidal. Perhaps it should be filtered to get out particulate matter but I have never seen this to be a problem. The pH of this has always turned out to be 7.4. My nurse discovered recently that if you do not shake the mixture to make it go into solution until after you

refrigerate it and are ready to use it that the solution is less yellow. I presume that this is good because sodium ascorbate is clear and dehydroascorbate is yellow. The made up solutions are always a little yellow but refrigeration before mixing results in a far less yellow mixture.

Preparation of the IV Bottle

I recommend that the above stock bottle solution be added to sterile water for injection such that 30 Gms (60 cc) to 60 Gms (120 cc) is added to a quantity of sterile water sufficient to make 500 cc of the final solution to be injected IV. For patients with chemical or other sensitivity reactions, it is best that the sterile water be obtained in glass IV bottles rather than the plastic IV bottles.

IM Injections

IM injection material for infants is made from the stock solution diluted 50% in water giving a 25% solution. Generally, the size of the injection can be 2 cc in each buttocks. Ice may be applied if it hurts too much. This may be given every hour or so, frequently enough to bring the fever or other symptoms of excessive free radicals down rapidly.

General Comments

I have not had any trouble with these solutions. I hear all sorts of weird stories from patients who have gotten ascorbate elsewhere. I do not know if it is an acid problem (because ascorbic acid was used rather than sodium ascorbate) or whether some colleges get carried away with what other things they add to the intravenous solutions.

I think that there may be, at times minor troubles with commercially prepared solutions because of the following. I understand that the U. S. Pharmacopeia specifies that the solutions be made from ascorbic acid and then buffered with sodium hydroxide or sodium bicarbonate to a pH between 3.5 and 7.0. I worry that 60 grams of ascorbate at a pH of 3.5 is too acid. I know that Klenner (the first physician who used high dose intravenous ascorbate by vein) also made his solutions from sodium ascorbate powder despite the fact that he referred to the solutions in his papers as ascorbic acid. (This fact comes from discussions with Annie Klenner, Fred Klenner's wife and nurse.).

I watch patients for hypocalcemia (although I have not seen it), hypoglycemia (I encourage patients to eat while taking the IV), and dehydration (I encourage water and slow the IV down.) I also see headaches afterward but not so much since I have been emphasizing the continuing high doses of oral ascorbic acid as soon as the IV is over. Actually I give oral ascorbic acid while the IV is going to get a double effect. Bowel tolerance goes up while the IV is running but one has to be careful to stop giving oral C about an hour before the IV stops or else you may get diarrhea as soon as the IV stops. The oral ascorbic acid is then started again 1/2 to 1 hour after the IVC stops.

Subscribe to the Orthomolecular Mailing List, for doctors, biochemists, and allied professionals. -- Write to Webmaster@mall-net.com.

Copyright (C) 1996, Robert F. Cathcart, M.D.. Permission granted to distribute as long as distributed intact and with this attribution.

Robert F. Cathcart, M.D.

650-949-2822



Publications by Robert F. Cathcart, M.D.

For your web space and web photography needs, contact webmaster@mall-net.com

Mall-Net